



SUPREMBIONic

Baja Ini Berasaskan Mikrob Aktif

Tidak mengandungi Mikrob Patogen, Salmonella Sp, As, Pb, Cd

Crops/Jenis Tanaman	Application Rate / Kadar Baja	Method of Application / Kaedah Pembajaan
Rice and Corn / Padi dan Jagung	<u>6 Liter untuk 1 hektar</u> a) 2 liter SUPREMBIONic + 200 liter air b) 1liter SUPREMBIONic + 100 liter air c) 2 liter SUPREMBIONic + 200 liter air d) 1 liter SUPREMBIONic + 100 liter air	<u>Kaedah :</u> a) Sembur 3 hari sebelum menanam benih padi / jagung b) Sembur hari ke-10 selepas ditanam c) Sembur hari ke-30 selepas ditanam d) Sembur semasa batang mula keluaran padi (primodial period of the rice)
Potato, Tomato, Pepper, Chilli, Soybeans, Peanuts / Ubi Kentang, Tomato, Kacang Soya, Kacang	<u>6 Liter untuk 1 hektar</u> a) 2 liter SUPREMBIONic + 100 liter air b) 2 liter SUPREMBIONic + 100 liter air c) 2 liter SUPREMBIONic + 100 liter air	<u>Kaedah :</u> a) Sembur 3 hari sebelum menanam benih b) Sembur hari ke-20 selepas ditanam c) Sembur hari ke-40 selepas ditanam
Vegetables / Sayur-sayuran	<u>6 Liter untuk 1 hektar</u> a) 2 liter SUPREMBIONic + 100 liter air b) 2 liter SUPREMBIONic + 100 liter air c) 2 liter SUPREMBIONic + 100 liter air	<u>Kaedah :</u> a) Sembur 3 hari sebelum menanam benih b) Sembur hari ke-20 selepas ditanam c) Sembur hari ke-40 selepas ditanam
Tobacco and Banana / Tembakau dan Pisang	<u>6 - 8 Liter untuk 1 hektar</u> a) 1 liter SUPREMBIONic + 150 liter air b) 1 liter SUPREMBIONic + 150 liter air	<u>Kaedah :</u> a) Sembur 3 hari sebelum menanam benih b) Sembur setiap 2 minggu

Ingredient:	Azotobacter Sp	$2.3 \times 10^{7-10^8}$	Azospirillum Sp	$3.0 \times 10^{7-10^8}$	Mikrob Pelarut Fosfat	$3.5 \times 10^{7-10^8}$
	Lactobacillus Sp	$2.0 \times 10^{7-10^8}$	Mikrob Selulotik	$2.0 \times 10^{7-10^8}$		

Berasaskan Teknologi Mesra Alam, BIO DEGRADABLE !!
Applied for both Edible Plants and Forest Floras

Hubungi :
 MUHAMMAD AFFAN
 014-9100623